

Post operative instructions for Excision of Vulvar Lesion

Pain

Pain is usually mild to moderate. If discomfort occurs, use acetaminophen (such as Tylenol) or ibuprofen (such as Advil) as directed. Avoid aspirin. A stronger pain medication (a narcotic) may be prescribed. Keeping ice on the area for 12 -- 24 hours helps reduce pain and swelling as well. Other medications you were taking before the surgery are generally restarted after surgery, but please review your medication

instructions and ask if you have any questions	about this.
Diet/Activity	
There are no dietary restrictions. After reco	overy from anesthesia is complete, there are no activity Yound Care". You may return to work/school in
Wound Care	
	eful wound care is important for fast and proper healing. Follow
Keep the area dry for 24 hours. If you minimize the moisture on the wound.	ou use ice to reduce pain, place the ice in a baggie to
	warm water several times per day if possible. This promotes
wound healing and relieves discomfort.	
Avoid intercourse for weeks	
Take one Colace stool softener (100	mg) twice per day for weeks
Things to watch and call for: If you develop a fever greater than 100.0 deg If you notice pus or foul smelling drainage fro If you have trouble urinating, burning when yo Please call if you have any concerns or question	om your wound. ou urinate, or feel the need to urinate very often.
Return or a post-operative visit in	
Call the office for this appointment. Phone numbers: Bel Air office: 443-643-4300,	Havre de Grace Office: 410-939-3121
Your Signature:	Date:
Physician Signature:	Date:
Nurse Signature:	Date:
	Bel Air

520 Upper Chesapeake Drive, Suite 301 • Bel Air, MD 21014 • Phone: 443.643.4300 • Fax: 443.643.4303

Havre de Grace 308 North Union Ave • Havre de Grace, MD 21078 • Phone: 410.939.3121 • Fax: 410.939.9411