



LAPAROSCOPICALLY ASSISTED VAGINAL HYSTERECTOMY
with BILATERAL SALPINGO-OOPHORECTOMY
Addendum to the authorization for surgical treatment

Complications and consequences have been discussed. I understand that the risks include, but are not limited to, the following:

- Pain (immediately afterwards, delayed, long term)
- Excessive bleeding, possibly requiring blood transfusion (and its risks of infection or transfusion reaction)
- Surgical infection
- Damage to abdominal/pelvic organs such as bladder, ureters, bowel, blood vessels, nerves possibly requiring further surgery or therapy at the time of the operation or at a later date
- Formation of blood clots in the legs and subsequent risk of pulmonary embolism
- Wound breakdown and bleeding in the vagina
- Formation of internal scar tissue (adhesions) causing pain, bowel obstruction, or difficulties with future surgeries
- Need for an abdominal incision to perform or complete the operation
- Risk of anesthesia
- Failure of procedure to cure the problem
- Urinary complications such as fistula formation, incontinence, or retention
- Onset of menopause after surgery, need to consider therapy for these symptoms and issues
- Prolapse of the vagina
- Other:

I acknowledge that I have read and fully understand the above complications and risks, and that all of my questions have been answered to my satisfaction.

Signature

Date

Witness

Date

Bel Air
520 Upper Chesapeake Drive, Suite 301 • Bel Air, MD 21014 • Phone: 443.643.4300 • Fax: 443.643.4303

Havre de Grace
308 North Union Ave • Havre de Grace, MD 21078 • Phone: 410.939.3121 • Fax: 410.939.9411